



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A SOLUTION OF EQUATIONS BY STANDARD CURVES.

By R. C. COLWELL.

If the general quadratic is expressed in the form

$$x^2 - kx + l = 0$$

$a + b = k$ can be represented by a straight line which cuts the




FIG. 1.

Solution: $x^2 - 5x + 3 = 0$, at R, S .

Solution: $x^2 - 2x + 3 = 0$, $OP + P'Q$.

hyperbola $ab = l$ in two points the co-ordinates of either of which are a and b , the roots of the quadratic.

Example (see Fig. 1) :

If $x^2 - 5x + 3 = 0$,

The line $a + b = 5$ cuts the hyperbola $ab = 3$ in the point R , S giving the solution $x = 4.3$ or $.7$.

As the value of $k^2 - 4l$ becomes smaller, l remaining constant, the line MN slides toward the origin, becoming tangent to the hyperbola at $k^2 - 4l = 0$, and not cutting the hyperbola at all when $k^2 - 4l < 0$. The roots of the equation are then complex and are determined by a point on the ellipse whose major axis is the major axis of the hyperbola and whose semi-minor axis equals $5l$. The real part of the complex root is evidently $k/2$, which is measured from the origin along the x axis. At the point thus found, a perpendicular is erected to cut the major




FIG. 2.

Solution: $x^2 + 4x - 5 = 0$, real root at A .

Imaginary roots, $OC \pm i \sqrt{BC}$.

axis of the ellipse. The perpendicular to the major axis from the point thus obtained cuts the ellipse and its length is the coefficient of i in the imaginary part of the root.

Example (Fig. 1):

If $x^3 - 2x + 3 = 0$,

The real part of the root is $OP = 1$;

The imaginary part is $P'Qi = 1.4i$;

The complete root $x = 1 \pm 1.4i$.

In the reduced cubic equation

$$x^3 + px + q = 0.$$

If a is one of the roots and c the product of the other two, then $a^2 - c = -p$ and $ac = -q$. Now $a^2 = c = -p$ is a parabola and $ac = -q$ is a hyperbola, and their points of intersec-




FIG. 3.

Solution: $x^3 - 3x - 2 = 0$. Roots at A and B .

tion will give the values of a and c , and therefore the roots.

Example (Fig. 2):

To solve $x^3 - 3x - 2 = 0$.

$a^2 - c = 3$. Therefore the vertex of the parabola $a^2 - c$ is at -3 on the y axis. It then cuts the hyperbola $ac = 2$ in the

point whose abscissa is 2 and touches the curve at the point whose abscissa is -1 . Therefore $a=2$, $c=1$, or $a=-1$, $c=-2$, from which the solution 2, -1 , -1 follows at once. If there is only one real root the parabola will cut the hyperbola in only one point; but the quadratic of the other two roots can be written at once and solved by the methods given before, or by the following method.

Example: $x^3 + 4x - 5 = 0$. (See Fig. 3.)

The parabola $a^2 - c = -4$ cuts the hyperbola $ac = 5$ in the point 1, 5. Therefore the real root is 1.

The quadratic of the remaining roots is

$$x^2 + x + 5 = 0.$$

This is solved by moving the major axis of the parabola, $a^2 = c$, which has been cut out, to the line $x = -\frac{1}{2}$ and letting it cut the y axis at 5. The ordinate of the vertex is the square of the coefficient of i .

The solution is then $-.5 \pm i\sqrt{4.7} = -.5 \pm 2.18i$.

The solution of a cubic in the reduced form

$$x^3 + (c - a^2)x - ac = 0$$

depends upon ac , and a^2 . The product ac may be found on the D scale of a slide rule with a and a^2 on the C and B scales respectively. It is therefore possible to solve a cubic in the above form upon a slide rule.*

GENEVA COLLEGE,
BEAVER FALLS, PA.

* Runge, "Graphical Methods."